

## SARS-CoV-2 (2019-nCoV) Spike RBD-Fc Recombinant Protein

Catalog_no :	AD-PD400024
Category :	冠状病毒产品
Size :	100µg/1mg
Specificity :	2019-nCoV
Source :	HEK293 Cells
Storage_stability :	Samples are stable for up to twelve months from date of receipt at -20°C to -80°C Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Molecular Weight :	The recombinant SARS-CoV-2 (2019-nCoV) Spike Protein (RBD, Fc Tag) consists of 461 amino acids and predicts a molecular mass of 51.8 kDa.
Background :	The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that 2019-nCoV can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.
缓冲液 :	Lyophilized from sterile PBS, pH 7.4. Please contact us for any concerns or special requirements. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information
运输及保存条件 :	In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice unless customers require otherwise
classification_1 :	coronavirus spike Protein, 2019-nCoV; cov spike Protein, 2019-nCoV; ncov RBD Protein, 2019-nCoV; ncov s1 Protein, 2019-nCoV; ncov s2 Protein, 2019-nCoV; ncov spike Protein, 2019-nCoV; NCP-CoV RBD Protein, 2019-nCoV; NCP-CoV s1 Protein, 2019-nCoV; NCP
purity :	> 95 % as determined by SDS-PAGE. > 95 % as determined by SEC-HPLC.

reference : 1.Shen S, et al. (2007) Expression, glycosylation, and modification of the spike (S) glycoprotein of SARS CoV. Methods Mol Biol. 379: 127-35. 2.Du L, et al. (2009) The spike protein of SARS-CoV--a target for vaccine and therapeutic development. Nat R

---

内毒素 : < 1.0 EU per  $\mu\text{g}$  protein as determined by the LAL method.

---

生物活性 : Measured by its binding ability in a functional ELISA. Immobilized human ACE2 protein (His tag) at  $2\mu\text{g}/\text{mL}$  ( $100\mu\text{L}/\text{well}$ ) can bind SARS-CoV-2 (2019-nCoV) Spike Protein (RBD, Fc Tag)(40592-V02H), the EC50 of 2019-nCoV Spike Protein (RBD, Fc Tag) is 2-6 n

---