

## PSMA5 Antibody (Ascites)

Catalog_no :	AB0711
Applications :	WB
Reactivity :	H
Category :	抗原抗体
Size :	100 $\mu$ L/50 $\mu$ L
Immunogen :	HUMAN
Specificity :	Purified His-tagged PSMA5 protein(Fragment) was used to produced this monoclonal antibody.
Dilution :	WB,1:1000~6400;
Purification :	Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.
Other_name :	Proteasome subunit alpha type-5, Macropain zeta chain, Multicatalytic endopeptidase complex zeta chain, Proteasome zeta chain, PSMA5
Isotype :	IgG1
Background :	The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit.
reference :	Kottgen, A., et al. Nat. Genet. 42(5):376-384(2010) Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007) Olsen, J.V., et al. Cell 127(3):635-648(2006) Beausoleil, S.A., et al. Nat. Biotechnol. 24(10):1285-1292(2006) Hirano, Y., et al. Natur