

TLL1 Antibody (C-term)

| | |
|----------------|--|
| Catalog_no : | AB0825 |
| Applications : | WB, IHC-P |
| Reactivity : | H |
| Category : | 抗原抗体 |
| Size : | 100μL/50μL |
| Immunogen : | HUMAN:874-901 |
| Specificity : | This TLL1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 874-901 amino acids from the C-terminal region of human TLL1. |
| Dilution : | IHC-P,1:50~100;WB,1:1000; |
| Purification : | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Other_name : | Tolloid-like protein 1, 3424-, TLL1, TLL |
| Isotype : | Rabbit Ig |
| Background : | This gene encodes an astacin-like zinc-dependent metalloprotease and is a subfamily member of the metzincin family. A similar protein in mice is required during heart development and specifically processes procollagen C-propeptides and chordin at similar cleavage sites. |
| reference : | Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) ; Berry, R., et al. FEBS Lett. 584(4):657-661(2010) Sabirzhanova, I.B., et al. Biochem. Biophys. Res. Commun. 389(2):338-342(2009) Stanczak, P., et al. Eur. J. Hum. Genet. 17(3):344-351(2009) Mac |