

## IER3 Antibody (N-term)

Catalog\_no: AB0961

Applications: WB

Reactivity: H, M

Category: 抗原抗体

Size:  $100\mu L/50\mu L$ 

Immunogen: HUMAN:38-65

Specificity: This IER3 antibody is generated from rabbits immunized with a KLH conjugated

synthetic peptide between 38-65 amino acids from the N-terminal region of human

IER3.

Dilution: WB,1:2000;WB,1:1000;WB,1:1000;

Purification: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This

antibody is purified through a protein A column, followed by peptide affinity

purification.

Other\_name: Radiation-inducible immediate-early gene IEX-1, Differentiation-dependent gene 2

protein, Protein DIF-2, Immediate early protein GLY96, Immediate early response 3

protein, PACAP-responsive gene 1 protein, Protein PRG1, IER3, DIF2, IEX1, PRG1

Isotype: Rabbit Ig

Background: This gene functions in the protection of cells from Fas- or tumor necrosis factor type

alpha-induced apoptosis. Partially degraded and unspliced transcripts are found after virus infection in vitro, but these transcripts are not found in vivo and do not generate a

valid protein.

reference: Shahid, M., et al. Hypertension 56(4):705-712(2010) Ucisik-Akkaya, E., et al. Mol. Hum.

Reprod. 16(10):770-777(2010) Fellay, I., et al. PLoS Genet. 5 (12), E1000791 (2009): Do,

T.N., et al. Cancer Genet. Cytogenet. 195(1):31-36(2009) Barcellos, L.F.