

## TAF8 Antibody (Center)

Catalog_no :	AB1180
Applications :	WB
Reactivity :	H
Category :	抗原抗体
Size :	100μL/50μL
Immunogen :	HUMAN:151-178
Specificity :	This TAF8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 151-178 amino acids from the Central region of human TAF8.
Dilution :	WB,1:1000;IHC-P,1:10~50;
Purification :	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Other_name :	Transcription initiation factor TFIID subunit 8, Protein taube nuss, TBP-associated factor 43 kDa, TBP-associated factor 8, Transcription initiation factor TFIID 43 kDa subunit, TAFII-43, TAFII43, hTAFII43, TAF8, TAFII43, TBN
Isotype :	Rabbit Ig
Background :	This gene encodes one of several TATA-binding protein (TBP)-associated factors (TAFs), which are integral subunits of the general transcription factor complex TFIID. TFIID recognizes the core promoter of many genes and nucleates the assembly of a transcription preinitiation complex containing RNA polymerase II and other initiation factors. The protein encoded by this gene contains an H4-like histone fold domain, and interacts with several subunits of TFIID including TBP and the histone-fold protein TAF10. Alternatively spliced transcript variants have been described, but their biological validity has not been determined. [provided by RefSeq].
reference :	Ganesh, S.K., et al. Nat. Genet. 41(11):1191-1198(2009) Soranzo, N., et al. Nat. Genet. 41(11):1182-1190(2009) Chapuis, J., et al. Mol. Psychiatry 14(11):1004-1016(2009) Soutoglou, E., et al. Mol. Cell. Biol. 25(10):4092-4104(2005) Guermah, M., et al