



## CTAG1A Antibody (N-term)

|                |  |
|----------------|--|
| Catalog_no :   | AB1194   |
| Applications : | WB   |
| Reactivity :   | H  |
| Category :     | 抗原抗体   |
| Size :         | 100μL/50μL   |
| Immunogen :    | HUMAN:1-30   |
| Specificity :  | This CTAG1A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human CTAG1A.  |
| Dilution :     | IHC-P,1:250;WB,1:1000;WB,1:2000;   |
| Purification : | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.   |
| Other_name :   | Cancer/testis antigen 1, Autoimmunogenic cancer/testis antigen NY-ESO-1, Cancer/testis antigen 61, CT61, L antigen family member 2, LAGE-2, CTAG1A, CTAG, CTAG1, ESO1, LAGE2, LAGE2A   |
| Isotype :      | Rabbit Ig  |
| Background :   | CTAG1A is an identical copy of the CTAG1B gene (MIM 300156) (Aradhya et al., 2001 [PubMed 11709543]).[supplied by OMIM].   |
| reference :    | Yoshida, N., et al. Int. J. Oncol. 28(5):1089-1098(2006) Cho, H.J., et al. Cancer Immun. 6, 12 (2006) : Ross, M.T., et al. Nature 434(7031):325-337(2005) Scanlan, M.J., et al. Immunol. Rev. 188, 22-32 (2002) : Satie, A.P., et al. Lab. Invest. 82(6):7 |