

TUBA3C Antibody (C-term)

| | |
|----------------|---|
| Catalog_no : | AB1198 |
| Applications : | WB |
| Reactivity : | H |
| Category : | 抗原抗体 |
| Size : | 100 μ L/50 μ L |
| Immunogen : | HUMAN:413-441 |
| Specificity : | This TUBA3C antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 413-441 amino acids from the C-terminal region of human TUBA3C. |
| Dilution : | WB,1:1000; |
| Purification : | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Other_name : | Tubulin alpha-3C/D chain, Alpha-tubulin 2, Alpha-tubulin 3C/D, Tubulin alpha-2 chain, TUBA3C, TUBA2 |
| Isotype : | Rabbit Ig |
| Background : | Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain. |
| reference : | Lamesch, P., et al. Genomics 89(3):307-315(2007) Villasante, A., et al. Mol. Cell. Biol. 6(7):2409-2419(1986) |