

CIR Antibody (N-term)

Catalog_no :	AB1299
Applications :	WB
Reactivity :	H
Category :	抗原抗体
Size :	100 μ L/50 μ L
Immunogen :	HUMAN:60-88
Specificity :	This CIR antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 60-88 amino acids from the N-terminal region of human CIR.
Dilution :	WB,1:1000;
Purification :	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Other_name :	Corepressor interacting with RBPJ 1, CBF1-interacting corepressor, Receptin, CIR1, CIR
Isotype :	Rabbit Ig
Background :	CIR may modulate splice site selection during alternative splicing of pre-mRNAs. It regulates transcription and acts as corepressor for RBPSUH. It recruits RBPSUH to the Sin3-histone deacetylase complex (HDAC) and is required for RBPSUH-mediated repression of transcription.
reference :	Olsen,J.V., Cell 127 (3), 635-648 (2006) Maita,H., Exp. Cell Res. 303 (2), 375-387 (2005) Zhou,S., Mol. Cell. Biol. 21 (18), 6222-6232 (2001)