

NAPB Antibody (N-term)

| | |
|----------------|--|
| Catalog_no : | AB1308 |
| Applications : | WB |
| Reactivity : | H |
| Category : | 抗原抗体 |
| Size : | 100μL/50μL |
| Immunogen : | HUMAN:13-40 |
| Specificity : | This NAPB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 13-40 amino acids from the N-terminal region of human NAPB. |
| Dilution : | WB,1:1000; |
| Purification : | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Other_name : | Beta-soluble NSF attachment protein, SNAP-beta, N-ethylmaleimide-sensitive factor attachment protein beta, NAPB, SNAPB |
| Isotype : | Rabbit Ig |
| Background : | NAPB is required for vesicular transport between the endoplasmic reticulum and the Golgi apparatus (By similarity). |
| reference : | Lamesch, P., et al. Genomics 89(3):307-315(2007) Prekeris, R., et al. J. Cell Biol. 143(4):957-971(1998) Osten, P., et al. Neuron 21(1):99-110(1998) Whiteheart, S.W., et al. Nature 362(6418):353-355(1993) Wilson, D.W., et al. J. Cell Biol. 117(3):531 |