

NFIC Antibody (Center)

Catalog_no :	AB1465
Applications :	WB
Reactivity :	H
Category :	抗原抗体
Size :	100μL/50μL
Immunogen :	HUMAN:180-208
Specificity :	This NFIC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 180-208 amino acids from the Central region of human NFIC.
Dilution :	WB,1:1000;IF,1:10~50;
Purification :	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Other_name :	Nuclear factor 1 C-type, NF1-C, Nuclear factor 1/C, CCAAT-box-binding transcription factor, CTF, Nuclear factor I/C, NF-I/C, NFI-C, TGGCA-binding protein, NFIC, NFI
Isotype :	Rabbit Ig
Background :	NFIC recognizes and binds the palindromic sequence 5'-TTGGCNNNNNGCCAA-3' present in viral and cellular promoters and in the origin of replication of adenovirus type 2. These proteins are individually capable of activating transcription and replication.
reference :	Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) ; Nilsson, J., et al. Cancer Res. 70(5):2020-2029(2010) Udelhoven, M., et al. J. Mol. Endocrinol. 44(2):99-113(2010) Plasari, G., et al. Mol. Cell. Biol. 29(22):6006-6017(2009) Riffel, A.K., et al