

## SNAPC2 Antibody (C-term)

Catalog_no :	AB1516
Applications :	WB
Reactivity :	H
Category :	抗原抗体
Size :	100μL/50μL
Immunogen :	HUMAN:278-307
Specificity :	This SNAPC2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 278-307 amino acids from the C-terminal region of human SNAPC2.
Dilution :	WB,1:1000;
Purification :	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Other_name :	snRNA-activating protein complex subunit 2, SNAPc subunit 2, Proximal sequence element-binding transcription factor subunit delta, PSE-binding factor subunit delta, PTF subunit delta, Small nuclear RNA-activating complex polypeptide 2, snRNA-activating protein complex 45 kDa subunit, SNAPc 45 kDa subunit, SNAPC2, SNAP45
Isotype :	Rabbit Ig
Background :	SNAPC2 is a subunit of the snRNA-activating protein complex which is associated with the TATA box-binding protein. The encoded protein is necessary for RNA polymerase II and III dependent small-nuclear RNA gene transcription. Alternative splicing results in multiple transcript variants. [provided by RefSeq].
reference :	Shanmugam, M., et al. J. Biol. Chem. 283(21):14845-14856(2008) Emran, F., et al. Gene 377, 96-108 (2006) : Hinkley, C.S., et al. J. Biol. Chem. 278(20):18649-18657(2003) Acierno, J.S. Jr., et al. Genomics 73(2):203-210(2001) Ma, B., et al. J. Biol. C