

BFAR Antibody (N-term)

| | |
|----------------|---|
| Catalog_no : | AB1550 |
| Applications : | WB |
| Reactivity : | H |
| Category : | 抗原抗体 |
| Size : | 100 μ L/50 μ L |
| Immunogen : | HUMAN:1-30 |
| Specificity : | This BFAR antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human BFAR. |
| Dilution : | WB,1:1000; |
| Purification : | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Other_name : | Bifunctional apoptosis regulator, RING finger protein 47, BFAR, BAR, RNF47 |
| Isotype : | Rabbit Ig |
| Background : | BFAR is a apoptosis regulator. Has anti-apoptotic activity, both for apoptosis triggered via death-receptors and via mitochondrial factors. |
| reference : | Liu, X., et al. Retina (Philadelphia, Pa.) (2010) In press : Chua, C.C., et al. Cardiovasc. Res. 81(1):20-27(2009) Roth, W., et al. Cell Death Differ. 10(10):1178-1187(2003) Zhang, H., et al. Proc. Natl. Acad. Sci. U.S.A. 97(6):2597-2602(2000) |