

HLF Antibody (C-term)

Catalog_no: AB1632

Applications: WB

Reactivity: H

Category: 抗原抗体

Size: $100\mu L/50\mu L$

Immunogen: HUMAN:204-232

Specificity: This HLF antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 204-232 amino acids from the C-terminal region of human HLF.

Dilution: WB,1:2000;

Purification: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This

antibody is purified through a protein A column, followed by peptide affinity

purification.

Other name: Hepatic leukemia factor, HLF

Isotype: Rabbit Ig

Background: This gene encodes a member of the proline and acidic-rich (PAR) protein family, a

subset of the bZIP transcription factors. The encoded protein forms homodimers or heterodimers with other PAR family members and binds sequence-specific promoter elements to activate transcription. Chromosomal translocations fusing portions of this gene with the E2A gene cause a subset of childhood B-lineage acute lymphoid leukemias. Alternatively spliced transcript variants have been described, but their

biological validity has not been determined.

reference: Hirose, K., et al. Blood 116(6):962-970(2010) Joslyn, G., et al. Alcohol. Clin. Exp. Res.

34(5):800-812(2010) Otto, T., et al. Endocrinology 149(5):2241-2250(2008) Inukai, T., et

al. Leukemia 21(2):288-296(2007) He, J., et al. Zhonghua Yi Xue Yi Chua