

PLEKHA8 Antibody (C-term)

| | |
|----------------|---|
| Catalog_no : | AB1765 |
| Applications : | WB |
| Reactivity : | H |
| Category : | 抗原抗体 |
| Size : | 100μL/50μL |
| Immunogen : | HUMAN:456-482 |
| Specificity : | This PLEKHA8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 456-482 amino acids from the C-terminal region of human PLEKHA8. |
| Dilution : | WB,1:1000; |
| Purification : | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Other_name : | Pleckstrin homology domain-containing family A member 8, PH domain-containing family A member 8, Phosphatidylinositol-four-phosphate adapter protein 2, FAPP-2, Phosphoinositol 4-phosphate adapter protein 2, hFAPP2, Serologically defined breast cancer antigen NY-BR-86, PLEKHA8, FAPP2 |
| Isotype : | Rabbit Ig |
| Background : | PLEKHA8 is involved in TGN-to-plasma membrane transport and in the formation of post-Golgi constitutive carriers. May play a role in ensuring the coordination of the budding and the fission reactions. |
| reference : | Tsao, D.A., et al. DNA Cell Biol. 29(6):285-293(2010) Cao, X., et al. Proc. Natl. Acad. Sci. U.S.A. 106(50):21121-21125(2009) Tritz, R., et al. Biochem. Biophys. Res. Commun. 383(2):167-171(2009) Yamaji, T., et al. IUBMB Life 60(8):511-518(2008) D'An |