

## MED19 Antibody (C-term)

Catalog_no :	AB1810
Applications :	WB, FC, IF
Reactivity :	H
Category :	抗原抗体
Size :	100μL/50μL
Immunogen :	HUMAN:161-188
Specificity :	This MED19 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 161-188 amino acids from the C-terminal region of human MED19.
Dilution :	WB,1:1000;
Purification :	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Other_name :	Mediator of RNA polymerase II transcription subunit 19, Lung cancer metastasis-related protein 1, Mediator complex subunit 19, MED19, LCMR1
Isotype :	Rabbit Ig
Background :	MED19 is a component of the Mediator complex, which is a coactivator for DNA-binding factors that activate transcription via RNA polymerase II (Sato et al., 2003 [PubMed 12584197]).[supplied by OMIM].
reference :	Ding, N., et al. J. Biol. Chem. 284(5):2648-2656(2009) Sato, S., et al. Mol. Cell 14(5):685-691(2004) Tomomori-Sato, C., et al. J. Biol. Chem. 279(7):5846-5851(2004) Sato, S., et al. J. Biol. Chem. 278(17):15123-15127(2003) Reddy, P.H., et al. Genomi