



FERMT2 Antibody (C-term)

Catalog_no :	AB1863
Applications :	WB
Reactivity :	H
Category :	抗原抗体
Size :	100μL/50μL
Immunogen :	HUMAN:641-669
Specificity :	This FERMT2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 641-669 amino acids from the C-terminal region of human FERMT2.
Dilution :	WB,1:1000;
Purification :	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Other_name :	Fermitin family homolog 2, Kindlin-2, Mitogen-inducible gene 2 protein, MIG-2, Pleckstrin homology domain-containing family C member 1, PH domain-containing family C member 1, FERMT2, KIND2, MIG2, PLEKHC1
Isotype :	Rabbit Ig
Background :	FERMT2 participates in the connection between ECM adhesion sites and the actin cytoskeleton and also in the orchestration of actin assembly and cell shape modulation. Recruits migfilin (FBLP1) protein to cell-ECM focal adhesion sites.
reference :	An, Z., et al. Int. J. Cancer 127(9):1999-2008(2010) Bledzka, K., et al. J. Biol. Chem. 285(40):30370-30374(2010) Lai-Cheong, J.E., et al. Int. J. Biochem. Cell Biol. 42(5):595-603(2010) Lai-Cheong, J.E., et al. J. Invest. Dermatol. 128(9):2156-2165(