

## NMI Antibody (N-term)

Catalog_no :	AB1911
Applications :	WB
Reactivity :	H
Category :	抗原抗体
Size :	100 $\mu$ L/50 $\mu$ L
Immunogen :	HUMAN:1-30
Specificity :	This NMI antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human NMI.
Dilution :	WB,1:1000;
Purification :	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Other_name :	N-myc-interactor, Nmi, N-myc and STAT interactor, NMI
Isotype :	Rabbit Ig
Background :	NMYC interactor (NMI) encodes a protein that interacts with NMYC and CMYC (two members of the oncogene Myc family), and other transcription factors containing a Zip, HLH, or HLH-Zip motif. The NMI protein also interacts with all STATs except STAT2 and augments STAT-mediated transcription in response to cytokines IL2 and IFN-gamma. The NMI mRNA has low expression levels in all human fetal and adult tissues tested except brain and has high expression in cancer cell line-myeloid leukemias. [provided by RefSeq].
reference :	Davila, S., et al. Genes Immun. 11(3):232-238(2010) Fillmore, R.A., et al. Int. J. Cancer 125(3):556-564(2009) Quaye, L., et al. Br. J. Cancer 100(6):993-1001(2009) Vega, A., et al. Gynecol. Oncol. 112(1):210-214(2009) Quaye, L., et al. Clin. Cancer