

AMHR2 Antibody (N-term)

Catalog_no: AB2476

Reactivity: H

Category: 抗原抗体

Size: $100\mu L/50\mu L$

Immunogen: HUMAN:13-42

Specificity: This AMHR2 antibody is generated from rabbits immunized with a KLH conjugated

synthetic peptide between 13-42 amino acids from the N-terminal region of human

AMHR2.

Dilution: WB,1:1000;

Purification: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This

antibody is purified through a protein G column, eluted with high and low pH buffers

and neutralized immediately, followed by dialysis against PBS.

Other_name: Anti-Muellerian hormone type-2 receptor, Anti-Muellerian hormone type II receptor,

AMH type II receptor, MIS type II receptor, MISRII, MRII, AMHR2, AMHR, MISR2

Isotype: Rabbit Ig

Background: The AMH receptor (AMHR or AMHR2) is a serine/threonine kinase with a single

transmembrane domain belonging to the family of type II receptors for TGF-beta-related proteins. Anti-Mullerian hormone (AMH) and its receptor are involved in the regression of Mullerian ducts in male fetuses. Male sex differentiation is mediated by 2 discrete hormones produced by the fetal testis. Testosterone, produced by Leydig cells, virilizes the external genitalia and promotes prostatic growth; anti-Mullerian hormone (AMH) results in regression of Mullerian ducts which would otherwise differentiate into

the uterus and fallopian tubes.

reference: Picard, J.Y., et al., J. Soc. Biol. 196(3):217-221 (2002). Teixeira, J., et al., Endocr. Rev.

22(5):657-674 (2001). Imbeaud, S., et al., Nat. Genet. 11(4):382-388 (1995). Visser, J.A., et

al., Biochem. Biophys. Res. Commun. 215(3):1029-1036 (1995). S