

PIST Antibody (C-term)

Catalog_no :	AB2489
Reactivity :	H, M
Category :	抗原抗体
Size :	100 μ L/50 μ L
Immunogen :	HUMAN:361-390
Specificity :	This PIST antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 361-390 amino acids from the C-terminal region of human PIST.
Dilution :	WB,1:1000;IHC-P,1:10~50;
Purification :	Purified rabbit polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS.
Other_name :	Golgi-associated PDZ and coiled-coil motif-containing protein, CFTR-associated ligand, Fused in glioblastoma, PDZ protein interacting specifically with TC10, PIST, GOPC, CAL, FIG
Isotype :	Rabbit Ig
Background :	PDZ domains contain approximately 90 amino acids and bind the extreme C terminus of proteins in a sequence-specific manner. PIST, a PDZ domain-containing Golgi protein, was discovered in a yeast two-hybrid system as a binding partner to Beclin-1, a Bcl-2-interacting protein homologous to the yeast autophagy gene apg6. Experiments with mutant PIST proteins lacking the PDZ domain showed that PIST interaction with Beclin-1 through its coiled-coil domain can modulate Beclin-1 activity and suggest that PIST interactions with other proteins through its PDZ domain may regulate the activity of PIST and Beclin-1.
reference :	Li,X., Protein Sci. 15 (9), 2149-2158 (2006) Ito,H., Biochem. J. 397 (3), 389-398 (2006) Wente,W., J. Biol. Chem. 280 (37), 32419-32425 (2005)