

## AGT Antibody (C-term)

Catalog_no :	AB2554
Reactivity :	H
Category :	抗原抗体
Size :	100 $\mu$ L/50 $\mu$ L
Immunogen :	HUMAN:418-450
Specificity :	This AGT antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 418-450 amino acids from the C-terminal region of human AGT.
Dilution :	WB,1:1000;IHC-P,1:10~50;
Purification :	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Other_name :	Angiotensinogen, Serpin A8, Angiotensin-1, Angiotensin 1-10, Angiotensin I, Ang I, Angiotensin-2, Angiotensin 1-8, Angiotensin II, Ang II, Angiotensin-3, Angiotensin 2-8, Angiotensin III, Ang III, Des-Asp[1]-angiotensin II, Angiotensin-4, Angiotensin 3-8, Angiotensin IV, Ang IV, Angiotensin 1-9, Angiotensin 1-7, Angiotensin 1-5, Angiotensin 1-4, AGT, SERPINA8
Isotype :	Rabbit Ig
Background :	AGT, pre-angiotensinogen or angiotensinogen precursor, is expressed in the liver and is cleaved by the enzyme renin in response to lowered blood pressure. The resulting product, angiotensin I, is then cleaved by angiotensin converting enzyme (ACE) to generate the physiologically active enzyme angiotensin II. The protein is involved in maintaining blood pressure and in the pathogenesis of essential hypertension and preeclampsia. Mutations in AGT gene are associated with susceptibility to essential hypertension, and can cause renal tubular dysgenesis, a severe disorder of renal tubular development. Defects in AGT gene have also been associated with non-familial structural atrial fibrillation, and inflammatory bowel disease.
reference :	Gurkan,A., Arch. Oral Biol. 54 (4), 337-344 (2009) Vickers,C., J. Biol. Chem. 277 (17), 14838-14843 (2002) Donoghue,M., Circ. Res. 87 (5), E1-E9 (2000)