

ASGR2 Antibody (N-term)

Catalog_no: AB3211

Reactivity: H

Category: 抗原抗体

Size: $100\mu L/50\mu L$

Immunogen: HUMAN

Specificity: This ASGR2 antibody is generated from a rabbit immunized with a KLH conjugated

synthetic peptide between 21-54 amino acids from the N-terminal region of human

ASGR2.

Dilution: WB,1:1000;

Other_name: Asialoglycoprotein receptor 2, ASGP-R 2, ASGPR 2, C-type lectin domain family 4

member H2, Hepatic lectin H2, HL-2, ASGR2, CLEC4H2

Isotype: Rabbit Ig

Background: Mediates the endocytosis of plasma glycoproteins to which the terminal sialic acid

residue on their complex carbohydrate moieties has been removed. The receptor recognizes terminal galactose and N-acetylgalactosamine units. After ligand binding to the receptor, the resulting complex is internalized and transported to a sorting

organelle, where receptor and ligand are disassociated. The receptor then returns to the

cell membrane surface.

reference: Spiess M., et al. Proc. Natl. Acad. Sci. U.S.A. 82:6465-6469(1985). Paietta E., et

al.Hepatology 15:395-402(1992). Zody M.C.,et al.Nature 440:1045-1049(2006). Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases. Pan H.,et

al.Genomic