

(Mouse) Ehmt2 Antibody (N-term)

Catalog_no :	AB3265
Reactivity :	H
Category :	抗原抗体
Size :	100 μ L/50 μ L
Immunogen :	HUMAN
Specificity :	This Mouse Ehmt2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 227-259 amino acids from the N-terminal region of Mouse Ehmt2.
Dilution :	WB,1:2000;
Other_name :	Histone-lysine N-methyltransferase EHMT2, 211-, Euchromatic histone-lysine N-methyltransferase 2, HLA-B-associated transcript 8, Histone H3-K9 methyltransferase 3, H3-K9-HMTase 3, Protein G9a, Ehmt2, Bat8, G9a, Ng36
Isotype :	Rabbit Ig
Background :	Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones. Also mediates monomethylation of 'Lys-56' of histone H3 (H3K56me1) in G1 phase, leading to promote interaction between histone H3 and PCNA and regulating DNA replication. Also weakly methylates 'Lys-27' of histone H3 (H3K27me). Also required for DNA methylation, the histone methyltransferase activity is not required for DNA methylation, suggesting that these 2 activities function independently. Probably targeted to histone H3 by different DNA-binding proteins like E2F6, MGA, MAX and/or DP1. May also methylate histone H1. In addition to the histone methyltransferase activity, also methylates non-histone proteins: mediates dimethylation of 'Lys- 373' of p53/TP53. Also methylates CDYL, WIZ, ACIN1, DNMT1, HDAC1, ERCC6, KLF12 and itself.
reference :	Tachibana M.,et al.Genes Dev. 16:1779-1791(2002). Xie T.,et al.Genome Res. 13:2621-2636(2003). Church D.M.,et al.PLoS Biol. 7:E1000112-E1000112(2009). Brown S.E.,et al.Mamm. Genome 12:916-924(2001). Tachibana M.,et al.J. Biol. Chem. 276:25309-25317(2001)