

(Mouse) Rnf2 Antibody (Center)

Catalog_no :	AB3360
Reactivity :	H, M, Rat
Category :	抗原抗体
Size :	100 μ L/50 μ L
Immunogen :	HUMAN
Specificity :	This Mouse Rnf2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 198-232 amino acids from the Central region of Mouse Rnf2.
Dilution :	WB,1:2000;WB,1:2000;
Other_name :	E3 ubiquitin-protein ligase RING2, 632-, RING finger protein 1B, RING1b, RING finger protein 2, Rnf2, DinG, Ring1b
Isotype :	Rabbit Ig
Background :	E3 ubiquitin-protein ligase that mediates monoubiquitination of 'Lys-119' of histone H2A (H2AK119Ub), thereby playing a central role in histone code and gene regulation. H2AK119Ub gives a specific tag for epigenetic transcriptional repression and participates in X chromosome inactivation of female mammals. May be involved in the initiation of both imprinted and random X inactivation. Essential component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones, rendering chromatin heritably changed in its expressibility. E3 ubiquitin-protein ligase activity is enhanced by BMI1/PCGF4. Acts as the main E3 ubiquitin ligase on histone H2A of the PRC1 complex, while RING1 may rather act as a modulator of RNF2/RING2 activity. Association to the chromosomal DNA is cell-cycle dependent. In resting B- and T-lymphocytes, interaction with AURKB leads to block its activity, thereby maintaining transcription in resting lymphocytes.
reference :	Schoorlemmer J.,et al.EMBO J. 16:5930-5942(1997). Carninci P.,et al.Science 309:1559-1563(2005). Ebert L.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases. Garcia E.,et al.EMBO J. 18:3404-3418(1999). Suzuki M.,et al.Development 129:4171-