

(Mouse) Trrap Antibody (C-term)

Catalog_no :	AB3380
Reactivity :	M
Category :	抗原抗体
Size :	100μL/50μL
Immunogen :	HUMAN
Specificity :	This mouse Trrap antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1703-1737 amino acids from the C-terminal region of mouse Trrap.
Dilution :	WB,1:2000;
Other_name :	Transformation/transcription domain-associated protein, Tra1 homolog, Trrap
Isotype :	Rabbit Ig
Background :	Adapter protein, which is found in various multiprotein chromatin complexes with histone acetyltransferase activity (HAT), which gives a specific tag for epigenetic transcription activation. Component of the NuA4 histone acetyltransferase complex which is responsible for acetylation of nucleosomal histones H4 and H2A. Plays a central role in MYC transcription activation, and also participates in cell transformation by MYC. Required for p53/TP53-, E2F1- and E2F4-mediated transcription activation. Probably acts by linking transcription factors such as E1A, MYC or E2F1 to HAT complexes such as STAGA thereby allowing transcription activation. Probably not required in the steps following histone acetylation in processes of transcription activation. May be required for the mitotic checkpoint and normal cell cycle progression. Component of a SWR1-like complex that specifically mediates the removal of histone H2A.Z/H2AFZ from the nucleosome.
reference :	Carninci P.,et al.Science 309:1559-1563(2005). Herceg Z.,et al.Nat. Genet. 29:206-211(2001).