

CHD3 Antibody (N-Term)

| | |
|---------------|--|
| Catalog_no : | AB3487 |
| Reactivity : | H |
| Category : | 抗原抗体 |
| Size : | 100μL/50μL |
| Immunogen : | HUMAN |
| Specificity : | This CHD3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 595-629 amino acids from human CHD3. |
| Dilution : | WB,1:1000-1:2000; |
| Other_name : | Chromodomain-helicase-DNA-binding protein 3, CHD-3, ATP-dependent helicase CHD3, Mi-2 autoantigen 240 kDa protein, Mi2-alpha, Zinc finger helicase, hZFH, CHD3 |
| Isotype : | Rabbit Ig |
| Background : | Component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin by deacetylating histones. Required for anchoring centrosomal pericentrin in both interphase and mitosis, for spindle organization and centrosome integrity. |
| reference : | Aubry F.,et al.Eur. J. Biochem. 254:558-564(1998). Zody M.C.,et al.Nature 440:1045-1049(2006). Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases. Woodage T.,et al.Proc. Natl. Acad. Sci. U.S.A. 94:11472-11477(1997). Ge Q.,et al. |