



CXXC5 Antibody (N-Term)

| | |
|---------------|--|
| Catalog_no : | AB3528 |
| Reactivity : | H |
| Category : | 抗原抗体 |
| Size : | 100μL/50μL |
| Immunogen : | HUMAN |
| Specificity : | This CXXC5 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 47-80 amino acids from human CXXC5. |
| Dilution : | WB,1:2000; |
| Other_name : | CXXC-type zinc finger protein 5, CF5, Putative MAPK-activating protein PM08, Putative NF-kappa-B-activating protein 102, Retinoid-inducible nuclear factor, RINF, CXXC5 |
| Isotype : | Rabbit Ig |
| Background : | May indirectly participate in activation of the NF- kappa-B and MAPK pathways. Acts as a mediator of BMP4-mediated modulation of canonical Wnt signaling activity in neural stem cells (By similarity). Required for DNA damage-induced ATM phosphorylation, p53 activation and cell cycle arrest. Involved in myelopoiesis. |
| reference : | Zhang M.,et al.Sci. China, Ser. C, Life Sci. 52:528-538(2009). Matsuda A.,et al.Oncogene 22:3307-3318(2003). Ota T.,et al.Nat. Genet. 36:40-45(2004). Zhou J.,et al.Submitted (JUL-2000) to the EMBL/GenBank/DDBJ databases. Schmutz J.,et al.Nature 431:2 |