

HIF1AN Antibody (C-term)

Catalog_no: AB3707

Reactivity: H

Category: 抗原抗体

Size: $100\mu L/50\mu L$

Immunogen: HUMAN

Specificity: This HIF1AN antibody is generated from a mouse immunized with a KLH conjugated

synthetic peptide between 1-349 amino acids from human HIF1AN.

Dilution: IF,1:25;WB,1:4000;

Other_name: Hypoxia-inducible factor 1-alpha inhibitor, 1.14.11.30, 1.14.11.n4, Factor inhibiting

HIF-1, FIH-1, Hypoxia-inducible factor asparagine hydroxylase, HIF1AN, FIH1

Isotype: IgG1,k

Background: Hydroxylates HIF-1 alpha at 'Asp-803' in the C-terminal transactivation domain (CAD).

Functions as an oxygen sensor and, under normoxic conditions, the hydroxylation prevents interaction of HIF-1 with transcriptional coactivators including Cbp/p300-interacting transactivator. Involved in transcriptional repression through interaction with HIF1A, VHL and histone deacetylases. Hydroxylates specific Asn residues within ankyrin repeat domains (ARD) of NFKB1, NFKBIA, NOTCH1, ASB4, PPP1R12A and several other ARD-containing proteins. Also hydroxylates Asp and His residues within ARDs of ANK1 and TNKS2, respectively. Negatively regulates NOTCH1 activity, accelerating myogenic differentiation. Positively regulates ASB4 activity, promoting vascular

differentiation.

reference: Mahon P.C.,et al.Genes Dev. 15:2675-2686(2001). Ota T.,et al.Nat. Genet. 36:40-45(2004).

Deloukas P., et al. Nature 429:375-381(2004). Mural R.J., et al. Submitted (SEP-2005) to the

EMBL/GenBank/DDBJ databases. Bechtel S.,et al.BMC Genomics 8:399-399(200