

## α-tubulin Monoclonal Antibody(8F11)

Catalog\_no: A1052

Applications: WB,IHC-p,IF,IP

Reactivity: Human, Rat, Mouse

Category: 抗原抗体

Size: 50µl/100µl

Immunogen: Recombinant Protein of Tubulin alpha-1A chain

Specificity: The antibody detects endogenousα-tubulin protein.

Formulation: PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol.

Source: Mouse

Dilution: WB: 1:500-10000 IP:1:200 IF 1:200 IHC 1:50-300

Purification: The antibody was affinity-purified from mouse ascites by affinity-chromatography using

specific immunogen.

Concentration: 1 mg/ml

Molecular

52KD

Weight:

Storage buffer: PBS,pH 7.4,containing 0.02% sodium azide as Preservative and 50% Glycerol

Store at: -20°C.Do not aliquot the antibody, For research use only


Background: Tubulin is one of several members of a small family of globular proteins. The tubulin

superfamilyincludes five distinct families, the alpha-, beta-, gamma-, delta-, and epsilon-

tubulins. The most commonmembers of the tubulin family are  $\alpha$ -tubulin an



## **Product Images**



Immunohistochemical analysis of paraffin-embedded Human-uterus-cancer tissue. 1,α-tubulin Monoclonal Antibody(8F11) was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room tempeRature, 30min). Negative control was used by secondary antibody only.