




Histone H3 Monoclonal Antibody(1G1)

| | |
|--------------------|--|
| Catalog_no : | A1055 |
| Applications : | WB,IHC-p,IF,IP |
| Reactivity : | Human,Mouse,Rat,Yeast |
| Category : | 抗原抗体 |
| Size : | 50μl/100μl |
| Immunogen : | Recombinant Protein of Histone H3 |
| Specificity : | The antibody detects endogenous Histone H3 protein. |
| Formulation : | PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol. |
| Source : | Mouse |
| Dilution : | WB: 1:2000-5000 IF:1:100-500 IP:1:200 IHC 1:50-300 |
| Purification : | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. |
| Concentration : | 1 mg/ml |
| Molecular Weight : | 15KD |
| Storage buffer : | PBS,pH 7.4,containing 0.02% sodium azide as Preservative and 50% Glycerol |
| Store at : | -20°C.Do not aliquot the antibody,For research use only |
| Background : | Histone H3 is one of the five main histone proteins involved in the structure of chromatin ineukaryotic cells. Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limitingDNA accessibility to the cellular machineries whic |

Product Images



Immunohistochemical analysis of paraffin-embedded Human-uterus tissue. 1,Histone H3 Monoclonal Antibody(1G1) was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.