



## Cleaved-MMP-10 (F99) Polyclonal Antibody

|                     |   |
|---------------------|---|
| Catalog_no :        | AC0059  |
| Applications :      | WB,ELISA  |
| Reactivity :        | Human   |
| Category :          | 抗原抗体  |
| Size :              | 100µg/50µg/20µg   |
| Gene_name :         | MMP10   |
| Protein_name :      | Stromelysin-2   |
| Humangene_id :      | <a href="#">4319</a>  |
| Humanswissprot_no : | <a href="#">P09238</a>  |
| Immunogen :         | The antiserum was produced against synthesized peptide derived from human MMP10. AA range:80-129  |
| Specificity :       | Cleaved-MMP-10 (F99) Polyclonal Antibody detects endogenous levels of fragment of activated MMP-10 protein resulting from cleavage adjacent to F99. |
| Formulation :       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source :            | Rabbit  |
| Dilution :          | Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.   |
| Purification :      | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.                               |
| Concentration :     | 1 mg/ml   |
| Storage_stability : | -20°C/1 year  |
| Msds :              | MSDS_Antibody.pdf   |
| Other_name :        | MMP10; STMY2; Stromelysin-2; SL-2; Matrix metalloproteinase-10; MMP-10; Transin-2   |
| Molecular Weight :  | 43KD  |

## Product Images



Western Blot analysis of various cells using Cleaved-MMP-10 (F99) Polyclonal Antibody diluted at 1 : 1000



Western Blot analysis of HepG2 cells using Cleaved-MMP-10 (F99) Polyclonal Antibody diluted at 1 : 1000