



Histone H3 (Mono Methyl Lys80) Polyclonal Antibody

Catalog_no :	<u>AH0005</u>
Applications :	<u>WB,ELISA</u>
Reactivity :	<u>Human,Mouse,Rat</u>
Category :	<u>抗原抗体</u>
Size :	<u>100µg/50µg/20µg</u>
Gene_name :	<u>HIST1H3A/HIST1H3/HIST1H3C/HIST1H3D/HIST1H3E/HIST1H3F/HIST1H3G/HIST1H3H/HIST1H3I/HIST1H3J/HIST2H3A/HIST2H3C/HIST2H3D/H3F3A/H3F3B/H3F3C</u>
Protein_name :	<u>Histone H3.1/Histone H3.2/Histone H3.3/Histone H3.3C</u>
Humangene_id :	<u>8350/8351/8352/8353/8354/8355/8356/8357/8358/8968/126961/333932/653604/3020/3021/440093</u>
Humanswissprot_no :	<u>P68431/Q71DI3/P84243/Q6NXT2</u>
Mousegene_id :	<u>319152/15077/15078/625328</u>
Ratgene_id :	<u>291159/100361558</u>
Ratswissprot_no :	<u>Q6LED0/P84245</u>
Immunogen :	<u>Synthesized peptide derived from human Histone H3 around the mono-methylation site of K80.</u>
Specificity :	<u>Mono-Methyl-Histone H3 (K80) Polyclonal Antibody detects endogenous levels of Histone H3 protein only when mono-methylated at K80.</u>
Formulation :	<u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u>
Source :	<u>Rabbit</u>
Dilution :	<u>Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.</u>
Purification :	<u>The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage_stability :	<u>-20°C/1 year</u>
Msds :	<u>MSDS_Antibody.pdf</u>
Other_name :	<u>HIST1H3A; H3FA; HIST1H3B; H3FL; HIST1H3C; H3FC; HIST1H3D; H3FB; HIST1H3E; H3FD;</u>




HIST1H3F; H3FI; HIST1H3G; H3FH; HIST1H3H; H3FK; HIST1H3I; H3FF; HIST1H3J; H3FJ;
Histone H3.1; Histone H3.1; Histone H3.2; Histone H3/m; Histone H3/o; H3F3A; H3.3A;
H3F3; PP781; H3F3B; H3.3B; Histone H3.3; H3F3C; Histone H3.3C; Histone H3.5

Molecular
Weight :

17KD

Product Images



Western Blot analysis of MCF-7 cells using Mono-Methyl-Histone H3 (K80) Polyclonal Antibody.. Secondary antibody was diluted at 1:20000