



HNF-3 α / β / γ (Acetyl Lys264/253/211) Polyclonal Antibody

| | |
|---------------------|--|
| Catalog_no : | AK0026 |
| Applications : | WB,ELISA |
| Reactivity : | Human,Mouse,Rat |
| Category : | 抗原抗体 |
| Size : | 100 μ g/50 μ g/20 μ g |
| Gene_name : | FOXA1/FOXA2/FOXA3 |
| Protein_name : | Hepatocyte nuclear factor 3-alpha/Hepatocyte nuclear factor 3-beta/Hepatocyte nuclear factor 3-gamma |
| Humangene_id : | 3169 |
| Humanswissprot_no : | P55317 |
| Mousegene_id : | 15375 |
| Mouseswissprot_no : | P35582 |
| Ratgene_id : | 25098 |
| Ratswissprot_no : | P23512 |
| Immunogen : | The antiserum was produced against synthesized Acetyl-peptide derived from human FOXA1 around the Acetylation site of Lys265. AA range:231-280 |
| Specificity : | Acetyl-HNF-3 α / β / γ (K264/253/211) Polyclonal Antibody detects endogenous levels of HNF-3 α / β / γ protein only when acetylated at K264/253/211. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Rabbit |
| Dilution : | Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage_stability : | -20°C/1 year |




Msd : [MSDS_Antibody.pdf](#)


Other_name : FOXA1; HNF3A; TCF3A; Hepatocyte nuclear factor 3-alpha; HNF-3-alpha; HNF-3A; Forkhead box protein A1; Transcription factor 3A; TCF-3A; FOXA2; HNF3B; TCF3B; Hepatocyte nuclear factor 3-beta; HNF-3-beta; HNF-3B; Forkhead box protein A2; Transcription factor 3B; TCF-3B; FOXA3; HNF3G; TCF3G; Hepatocyte nuclear factor 3-gamma; HNF-3-gamma; HNF-3G; Fork head-related protein FKH H3; Forkhead box protein A3; Transcription factor 3G; TCF-3G

Molecular Weight : 50KD

Product Images



Western Blot analysis of Jurkat cells using Acetyl-HNF-3 α / β / γ (K264/253/211) Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Western Blot analysis of JK cells using Acetyl-HNF-3 α / β / γ (K264/253/211) Polyclonal Antibody. Secondary antibody was diluted at 1:20000