

## Histone H2B (Acetyl Lys126) Polyclonal Antibody

Catalog_no :	<u>AK0027</u>
Applications :	<u>WB,ELISA</u>
Reactivity :	<u>Human,Mouse,Rat</u>
Category :	<u>抗原抗体</u>
Size :	<u>100µg/50µg/20µg</u>
Gene_name :	<u>HIST1H2BC</u>
Protein_name :	<u>Histone H2B type 1-A/Histone H2B type 1-B/Histone H2B type 1-C/E/F/G/I</u>
Humangene_id :	<u><a href="#">255626</a></u>
Humanswissprot_no :	<u><a href="#">Q96A08</a></u>
Mousegene_id :	<u><a href="#">319177</a></u>
Mouseswissprot_no :	<u><a href="#">P70696</a></u>
Ratgene_id :	<u><a href="#">24829</a></u>
Ratswissprot_no :	<u><a href="#">Q00729</a></u>
Immunogen :	<u>The antiserum was produced against synthesized Acetyl-peptide derived from human H2B around the Acetylation site of Lys126. AA range:78-127</u>
Specificity :	<u>Acetyl-Histone H2B (K126) Polyclonal Antibody detects endogenous levels of Histone H2B protein only when acetylated at K126.</u>
Formulation :	<u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u>
Source :	<u>Rabbit</u>
Dilution :	<u>Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.</u>
Purification :	<u>The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage_stability :	<u>-20°C/1 year</u>
Msds :	<u>MSDS_Antibody.pdf</u>




Other\_name : HIST1H2BA; TSH2B; Histone H2B type 1-A; Histone H2B, testis; Testis-specific histone H2B; HIST1H2BB; H2BFF; Histone H2B type 1-B; Histone H2B.1; Histone H2B.f; H2B/f; HIST1H2BC; H2BFL; HIST1H2BE; H2BFH; HIST1H2BF;H2BFG; HIST1H2BG; H2BFA; HIST1H2BI; H2BFK; Histone H2B type 1-C/E/F/G/I; Histone H2B.1 A; Histone H2B.a; H2B/a; Histone H2B.g; H2B/g; Histone H2B.h; H2B/h; Histone H2B.k; H2B/k; Histone H2B.l; H2B/l

---


Molecular Weight : 15KD

---

## Product Images



Western Blot analysis of A549 cells using Acetyl-Histone H2B (K126) Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Western Blot analysis of A549 cells using Acetyl-Histone H2B (K126) Polyclonal Antibody. Secondary antibody was diluted at 1:20000