

NFκB-p65 (Acetyl Lys221) Polyclonal Antibody


| | |
|---------------------|---|
| Catalog_no : | AK0065 |
| Applications : | WB,ELISA |
| Reactivity : | Human,Mouse,Rat |
| Category : | 抗原抗体 |
| Size : | 100μg/50μg/20μg |
| Gene_name : | RELA |
| Protein_name : | Transcription factor p65 |
| Humangene_id : | 5970 |
| Humanswissprot_no : | Q04206 |
| Mousegene_id : | 19697 |
| Mouseswissprot_no : | Q04207 |
| Ratgene_id : | |
| Ratswissprot_no : | |
| Immunogen : | Synthesized acetyl-peptide derived from the Internal region of human NFκB-p65 around the acetylation site of K221. |
| Specificity : | Acetyl-NFκB-p65 (K221) Polyclonal Antibody detects endogenous levels of NFκB-p65 protein only when acetylation at K221. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Rabbit |
| Dilution : | Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage_stability : | -20°C/1 year |
| Msds : | MSDS_Antibody.pdf |




Other_name : RELA; NFKB3; Transcription factor p65; Nuclear factor NF-kappa-B p65 subunit; Nuclear factor of kappa light polypeptide gene enhancer in B-cells 3

Molecular Weight : 65KD

Product Images



Western Blot analysis of HepG2-UV, mouse brain cells using Acetyl-NFκB-p65 (K221) Polyclonal Antibody. Antibody was diluted at 1:1000. Secondary antibody was diluted at 1:20000



Western Blot analysis of HEPG2-UV cells using Acetyl-NFκB-p65 (K221) Polyclonal Antibody diluted at 1 : 1000. Secondary antibody was diluted at 1:20000