



CD5 Monoclonal Antibody(10G8)

Catalog_no :	AM3071
Applications :	IHC-P,IF
Reactivity :	Human,Mouse,Rat
Category :	抗原抗体
Size :	100µg/50µg
Gene_name :	CD5
Protein_name :	T-cell surface glycoprotein CD5
Humangene_id	921
:	
Humanswissprot	P06127
_no :	
Mousegene_id :	12507
Mouseswissprot	P13379
_no :	
Ratgene_id :	54236
Ratswissprot_no	P51882
:	
Immunogen :	Synthetic Peptide of CD5
Specificity :	The antibody detects endogenous CD5 proteins.
Formulation :	PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol.
Source :	Mouse
Dilution :	IHC 1:200 IF 1:50-200
Purification :	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Storage_stability	-20°C/1 year
:	
Msds :	MSDS_Antibody.pdf
Other_name :	CD5; LEU1; T-cell surface glycoprotein CD5; Lymphocyte antigen T1/Leu-1; CD5


Product Images




Immunohistochemical analysis of paraffin-embedded Human-liver-cancer tissue. 1,CD5 Monoclonal Antibody(10G8) was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.



Immunohistochemical analysis of paraffin-embedded Rat-kidney tissue. 1,CD5 Monoclonal Antibody(10G8) was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.



Immunohistochemical analysis of paraffin-embedded Mouse-kidney tissue. 1,CD5 Monoclonal Antibody(10G8) was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.



A

B

C

Immunofluorescence analysis of Mouse-heart tissue. 1,CD5 Monoclonal Antibody(10G8)(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B

Immunohistochemical analysis of paraffin-embedded Human Tonsil Carcinoma using CD5 Mouse mAb diluted at 1:200.

