

## NFκB-p65 (phospho Ser468) Polyclonal Antibody

|                     |   |
|---------------------|---|
| Catalog_no :        | AP0188  |
| Applications :      | WB,IHC-p,IP,ELISA   |
| Reactivity :        | Human,Mouse,Rat   |
| Category :          | 抗原抗体  |
| Size :              | 100μg/50μg/20μg   |
| Gene_name :         | RELA  |
| Protein_name :      | Transcription factor p65  |
| Humangene_id :      | <a href="#">5970</a>  |
| Humanswissprot_no : | <a href="#">Q04206</a>  |
| Mousegene_id :      | <a href="#">19697</a>   |
| Mouseswissprot_no : | <a href="#">Q04207</a>  |
| Immunogen :         | The antiserum was produced against synthesized peptide derived from human NF-kappaB p65 around the phosphorylation site of Ser468. AA range:435-484             |
| Specificity :       | Phospho-NFκB-p65 (S468) Polyclonal Antibody detects endogenous levels of NFκB-p65 protein only when phosphorylated at S468.                                     |
| Formulation :       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source :            | Rabbit  |
| Dilution :          | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunoprecipitation: 2-5 ug/mg lysate. ELISA: 1/20000. Not yet tested in other applications. |
| Purification :      | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Concentration :     | 1 mg/ml   |
| Storage_stability : | -20°C/1 year  |
| Msds :              | <a href="#">MSDS_Antibody.pdf</a>   |
| Other_name :        | RELA; NFKB3; Transcription factor p65; Nuclear factor NF-kappa-B p65 subunit; Nuclear factor of kappa light polypeptide gene enhancer in B-cells 3              |



Molecular  
Weight :

60KD

---

## Product Images



Western Blot analysis of various cells using Phospho-NFκB-p65 (S468) Polyclonal Antibody diluted at 1 : 1000