

NFATc4 (phospho Ser676) Polyclonal Antibody

| | |
|-------------------|---|
| Catalog_no : | AP0313 |
| Applications : | WB,IHC-p,ELISA |
| Reactivity : | Human,Mouse |
| Category : | 抗原抗体 |
| Size : | 100µg/50µg/20µg |
| Gene_name : | NFATC4 |
| Protein_name : | Nuclear factor of activated T-cells cytoplasmic 4 |
| Humangene_id | 4776 |
| : | |
| Humanswissprot | Q14934 |
| _no : | |
| Mousegene_id : | 73181 |
| Mouseswissprot | Q8K120 |
| _no : | |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human NFAT3 around the phosphorylation site of Ser676. AA range:642-691 |
| Specificity : | Phospho-NFATc4 (S676) Polyclonal Antibody detects endogenous levels of NFATc4 protein only when phosphorylated at S676. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Rabbit |
| Dilution : | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications. |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage_stability | -20°C/1 year |
| : | |
| Msds : | MSDS_Antibody.pdf |
| Other_name : | NFATC4; NFAT3; Nuclear factor of activated T-cells; cytoplasmic 4; NF-ATc4; NFATc4; T-cell transcription factor NFAT3; NF-AT3 |



Molecular
Weight :

120KD
