

## NFκB-p100 (phospho Ser872) Polyclonal Antibody

|                     |  |
|---------------------|--|
| Catalog_no :        | AP0375   |
| Applications :      | WB,ELISA   |
| Reactivity :        | Human,Mouse  |
| Category :          | 抗原抗体   |
| Size :              | 100μg/50μg/20μg  |
| Gene_name :         | NFKB2  |
| Protein_name :      | Nuclear factor NF-kappa-B p100 subunit   |
| Humangene_id :      | <a href="#">4791</a>   |
| Humanswissprot_no : | <a href="#">Q00653</a>   |
| Mousegene_id :      | <a href="#">18034</a>  |
| Mouseswissprot_no : | <a href="#">Q9WTK5</a>   |
| Immunogen :         | The antiserum was produced against synthesized peptide derived from human NF-kappaB p100 around the phosphorylation site of Ser872. AA range:838-887   |
| Specificity :       | Phospho-NFκB-p100 (S872) Polyclonal Antibody detects endogenous levels of NFκB-p100 protein only when phosphorylated at S872.  |
| Formulation :       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source :            | Rabbit   |
| Dilution :          | Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.  |
| Purification :      | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| Concentration :     | 1 mg/ml  |
| Storage_stability : | -20°C/1 year   |
| Msds :              | MSDS_Antibody.pdf  |
| Other_name :        | NFKB2; LYT10; Nuclear factor NF-kappa-B p100 subunit; DNA-binding factor KBF2; H2TF1; Lymphocyte translocation chromosome 10 protein; Nuclear factor of kappa light polypeptide gene enhancer in B-cells 2; Oncogene Lyt-10; Lyt10 |



Molecular  
Weight :

96KD

---