

Synuclein- α (phospho Tyr136) Polyclonal Antibody

| | |
|---------------------|--|
| Catalog_no : | AP0521 |
| Applications : | WB,ELISA |
| Reactivity : | Human,Mouse,Rat |
| Category : | 抗原抗体 |
| Size : | 100 μ g/50 μ g/20 μ g |
| Gene_name : | SNCA |
| Protein_name : | Alpha-synuclein |
| Humangene_id : | 6622 |
| Humanswissprot_no : | P37840 |
| Mousegene_id : | 20617 |
| Mouseswissprot_no : | O55042 |
| Ratgene_id : | 29219 |
| Ratswissprot_no : | P37377 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human Synuclein-alpha around the phosphorylation site of Tyr136. AA range:91-140 |
| Specificity : | Phospho-Synuclein- α (Y136) Polyclonal Antibody detects endogenous levels of Synuclein- α protein only when phosphorylated at Y136. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Rabbit |
| Dilution : | Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications. |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage_stability : | -20°C/1 year |
| Msds : | MSDS_Antibody.pdf |



Other_name : SNCA; NACP; PARK1; Alpha-synuclein; Non-A beta component of AD amyloid; Non-A4 component of amyloid precursor; NACP

Molecular Weight : 20KD
