

Cytokeratin 18 (phospho Ser52) Polyclonal Antibody

Catalog_no: AP0547

Applications: WB,IHC-p,IF,ELISA

Reactivity: Human

Category: 抗原抗体

Size: 100μg/50μg/20μg

Gene_name: KRT18

Protein_name : Keratin type I cytoskeletal 18

Humangene_id 3875

Humanswissprot P05783

_no:

The antiserum was produced against synthesized peptide derived from human Keratin Immunogen:

18 around the phosphorylation site of Ser52. AA range:21-70

Phospho-Cytokeratin 18 (S52) Polyclonal Antibody detects endogenous levels of Specificity:

Cytokeratin 18 protein only when phosphorylated at S52.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Rabbit

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other

applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography

using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage_stability -20°C/1 year

Msds: MSDS_Antibody.pdf

48KD

KRT18; CYK18; PIG46; Keratin; type I cytoskeletal 18; Cell proliferation-inducing gene 46 Other_name:


protein; Cytokeratin-18; CK-18; Keratin-18; K18

Molecular


Weight:



Product Images



Western blot analysis of 3T3 293T using p-Cytokeratin 18 (S52) antibody. Antibody was diluted at 1:500



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.