

## Lck (phospho Tyr505) Polyclonal Antibody

|                     |   |
|---------------------|---|
| Catalog_no :        | AP0660  |
| Applications :      | WB,IHC-p,ELISA  |
| Reactivity :        | Human,Mouse,Rat   |
| Category :          | 抗原抗体  |
| Size :              | 100µg/50µg/20µg   |
| Gene_name :         | LCK   |
| Protein_name :      | Tyrosine-protein kinase Lck   |
| Humangene_id :      | <a href="#">3932</a>  |
| Humanswissprot_no : | <a href="#">P06239</a>  |
| Mousegene_id :      | <a href="#">16818</a>   |
| Mouseswissprot_no : | <a href="#">P06240</a>  |
| Ratgene_id :        | <a href="#">313050</a>  |
| Ratswissprot_no :   | <a href="#">Q01621</a>  |
| Immunogen :         | The antiserum was produced against synthesized peptide derived from human Lck around the phosphorylation site of Tyr505. AA range:460-509 |
| Specificity :       | Phospho-Lck (Y505) Polyclonal Antibody detects endogenous levels of Lck protein only when phosphorylated at Y505.                         |
| Formulation :       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source :            | Rabbit  |
| Dilution :          | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.                  |
| Purification :      | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.                     |
| Concentration :     | 1 mg/ml   |
| Storage_stability : | -20°C/1 year  |



---

|                    |  |
|--------------------|--|
| MsdS :             | <a href="#">MSDS_Antibody.pdf</a>  |
| Other_name :       | LCK; Tyrosine-protein kinase Lck; Leukocyte C-terminal Src kinase; LSK; Lymphocyte cell-specific protein-tyrosine kinase; Protein YT16; Proto-oncogene Lck; T cell-specific protein-tyrosine kinase; p56-LCK |
| Molecular Weight : | 56KD   |

---