

## RelB (phospho Ser552) Polyclonal Antibody

|                     |  |
|---------------------|--|
| Catalog_no :        | AP0844   |
| Applications :      | WB,IHC-p,ELISA   |
| Reactivity :        | Human,Mouse  |
| Category :          | 抗原抗体   |
| Size :              | 100µg/50µg/20µg  |
| Gene_name :         | RELB   |
| Protein_name :      | Transcription factor RelB  |
| Humangene_id :      | <a href="#">5971</a>   |
| Humanswissprot_no : | <a href="#">Q01201</a>   |
| Mousegene_id :      | <a href="#">19698</a>  |
| Mouseswissprot_no : | <a href="#">Q04863</a>   |
| Immunogen :         | The antiserum was produced against synthesized peptide derived from human RelB around the phosphorylation site of Ser552. AA range:530-579 |
| Specificity :       | Phospho-RelB (S552) Polyclonal Antibody detects endogenous levels of RelB protein only when phosphorylated at S552.                        |
| Formulation :       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source :            | Rabbit   |
| Dilution :          | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.                   |
| Purification :      | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.                      |
| Concentration :     | 1 mg/ml  |
| Storage_stability : | -20°C/1 year   |
| Msds :              | <a href="#">MSDS_Antibody.pdf</a>  |
| Other_name :        | RELB; Transcription factor RelB; I-Rel   |
| Molecular           | 62KD   |



Weight :

---