

NFATc1 (phospho Ser294) Polyclonal Antibody

| | |
|---------------------|--|
| Catalog_no : | AP0958 |
| Applications : | IHC-p,ELISA |
| Reactivity : | Human,Mouse |
| Category : | 抗原抗体 |
| Size : | 100µg/50µg/20µg |
| Gene_name : | NFATC1 |
| Protein_name : | Nuclear factor of activated T-cells cytoplasmic 1 |
| Humangene_id : | 4772 |
| Humanswissprot_no : | O95644 |
| Mouseswissprot_no : | O88942 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human NFAT2 around the phosphorylation site of Ser294. AA range:261-310 |
| Specificity : | Phospho-NFATc1 (S294) Polyclonal Antibody detects endogenous levels of NFATc1 protein only when phosphorylated at S294. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Rabbit |
| Dilution : | Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage_stability : | -20°C/1 year |
| Msd : | MSDS_Antibody.pdf |
| Other_name : | NFATC1; NFAT2; NFATC; Nuclear factor of activated T-cells; cytoplasmic 1; NF-ATc1; NFATc1; NFAT transcription complex cytosolic component; NF-ATc; NFATc |