



## FX Polyclonal Antibody

|                     |   |
|---------------------|---|
| Catalog_no :        | AT1802  |
| Applications :      | WB,ELISA  |
| Reactivity :        | Human,Mouse,Rat   |
| Category :          | 抗原抗体  |
| Size :              | 100µg/50µg/20µg   |
| Gene_name :         | TSTA3   |
| Protein_name :      | GDP-L-fucose synthase   |
| Humangene_id :      | <a href="#">7264</a>  |
| Humanswissprot_no : | <a href="#">Q13630</a>  |
| Mousegene_id :      | <a href="#">22122</a>   |
| Mouseswissprot_no : | <a href="#">P23591</a>  |
| Immunogen :         | The antiserum was produced against synthesized peptide derived from human TSTA3. AA range:221-270   |
| Specificity :       | FX Polyclonal Antibody detects endogenous levels of FX protein.   |
| Formulation :       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source :            | Rabbit  |
| Dilution :          | Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.   |
| Purification :      | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Concentration :     | 1 mg/ml   |
| Storage_stability : | -20°C/1 year  |
| Msds :              | <a href="#">MSDS_Antibody.pdf</a>   |
| Other_name :        | TSTA3; SDR4E1; GDP-L-fucose synthase; GDP-4-keto-6-deoxy-D-mannose-3; 5-epimerase-4-reductase; Protein FX; Red cell NADP(H)-binding protein; Short-chain dehydrogenase/reductase family 4E member 1 |
| Molecular           | 40KD  |



Weight :

---