


Hic-5 Polyclonal Antibody

| | |
|---------------------|--|
| Catalog_no : | AT2132 |
| Applications : | WB,IHC-p,IF,ELISA |
| Reactivity : | Human,Mouse,Rat |
| Category : | 抗原抗体 |
| Size : | 100µg/50µg/20µg |
| Gene_name : | TGFB1I1 |
| Protein_name : | Transforming growth factor beta-1-induced transcript 1 protein |
| Humangene_id : | 7041 |
| Humanswissprot_no : | O43294 |
| Mousegene_id : | 21804 |
| Mouseswissprot_no : | Q62219 |
| Ratgene_id : | 84574 |
| Ratswissprot_no : | Q99PD6 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human Hic-5. AA range:31-80 |
| Specificity : | Hic-5 Polyclonal Antibody detects endogenous levels of Hic-5 protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Rabbit |
| Dilution : | Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage_stability : | -20°C/1 year |
| Msds : | MSDS_Antibody.pdf |



Other_name : TGFB1I1; ARA55; Transforming growth factor beta-1-induced transcript 1 protein; Androgen receptor coactivator 55 kDa protein; Androgen receptor-associated protein of 55 kDa; Hydrogen peroxide-inducible clone 5 protein; Hic-5

Product Images



Western Blot analysis of NIH-3T3 cells using Hic-5
Polyclonal Antibody diluted at 1 : 1000