

ICAD Polyclonal Antibody


Catalog_no :	AT2268
Applications :	WB,IHC-p,IF,ELISA
Reactivity :	Human,Mouse,Rat
Category :	抗原抗体
Size :	100µg/50µg/20µg
Gene_name :	DFFA
Protein_name :	DNA fragmentation factor subunit alpha
Humangene_id :	1676
Humanswissprot_no :	O00273
Mousegene_id :	13347
Mouseswissprot_no :	O54786
Immunogen :	The antiserum was produced against synthesized peptide derived from human DFFA. AA range:151-200
Specificity :	ICAD Polyclonal Antibody detects endogenous levels of ICAD protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Rabbit
Dilution :	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage_stability :	-20°C/1 year
Msds :	MSDS_Antibody.pdf
Other_name :	DFFA; DFF1; DFF45; H13; DNA fragmentation factor subunit alpha; DNA fragmentation factor 45 kDa subunit; DFF-45; Inhibitor of CAD; ICAD




Molecular
Weight :

36KD

Product Images



Western Blot analysis of various cells using ICAD Polyclonal Antibody diluted at 1 : 1000



Western Blot analysis of 3T3 cells using ICAD Polyclonal Antibody diluted at 1 : 1000