

## NF-YC Polyclonal Antibody


Catalog_no :	<u>AT3092</u>
Applications :	<u>WB,IHC-p,ELISA</u>
Reactivity :	<u>Human,Mouse,Rat</u>
Category :	<u>抗原抗体</u>
Size :	<u>100µg/50µg/20µg</u>
Gene_name :	<u>NFYC</u>
Protein_name :	<u>Nuclear transcription factor Y subunit gamma</u>
Humangene_id	<u><a href="#">4802</a></u>
:	<u></u>
Humanswissprot	<u><a href="#">Q13952</a></u>
_no :	<u></u>
Mousegene_id :	<u><a href="#">18046</a></u>
Mouseswissprot	<u><a href="#">P70353</a></u>
_no :	<u></u>
Ratgene_id :	<u><a href="#">25337</a></u>
Ratswissprot_no	<u><a href="#">Q62725</a></u>
:	<u></u>
Immunogen :	<u>The antiserum was produced against synthesized peptide derived from human NFYC. AA range:1-50</u>
Specificity :	<u>NF-YC Polyclonal Antibody detects endogenous levels of NF-YC protein.</u>
Formulation :	<u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u>
Source :	<u>Rabbit</u>
Dilution :	<u>Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.</u>
Purification :	<u>The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage_stability	<u>-20°C/1 year</u>
:	<u></u>
Msds :	<u>MSDS_Antibody.pdf</u>




Other\_name : NFYC; Nuclear transcription factor Y subunit gamma; CAAT box DNA-binding protein subunit C; Nuclear transcription factor Y subunit C; NF-YC; Transactivator HSM-1/2

Molecular Weight : 50KD

## Product Images



Western Blot analysis of various cells using NF-YC Polyclonal Antibody diluted at 1 : 1000



Western Blot analysis of 293 cells using NF-YC Polyclonal Antibody diluted at 1 : 1000