





## Nop132 Polyclonal Antibody

Catalog_no :	AT3165
Applications :	WB,IHC-p,ELISA
Reactivity :	Human
Category :	抗原抗体
Size :	100µg/50µg/20µg
Gene_name :	NOL8
Protein_name :	Nucleolar protein 8
Humangene_id :	<a href="#">55035</a>
Humanswissprot_no :	<a href="#">Q76FK4</a>
Immunogen :	The antiserum was produced against synthesized peptide derived from human NOL8. AA range:1118-1167
Specificity :	Nop132 Polyclonal Antibody detects endogenous levels of Nop132 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Rabbit
Dilution :	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage_stability :	-20°C/1 year
Msds :	MSDS_Antibody.pdf
Other_name :	NOL8; C9orf34; NOP132; Nucleolar protein 8; Nucleolar protein Nop132
Molecular Weight :	130KD

## Product Images



Western Blot analysis of 3T3 cells using Nop132 Polyclonal Antibody diluted at 1 : 1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Western Blot analysis of NIH-3T3 cells using Nop132 Polyclonal Antibody diluted at 1 : 1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).