

## Oct-2 Polyclonal Antibody

|                     |  |
|---------------------|--|
| Catalog_no :        | AT3231   |
| Applications :      | WB,IHC-p,IF,ELISA  |
| Reactivity :        | Human,Mouse,Monkey   |
| Category :          | 抗原抗体   |
| Size :              | 100µg/50µg/20µg  |
| Gene_name :         | POU2F2   |
| Protein_name :      | POU domain class 2 transcription factor 2  |
| Humangene_id :      | <a href="#">5452</a>   |
| Humanswissprot_no : | <a href="#">P09086</a>   |
| Mousegene_id :      | <a href="#">18987</a>  |
| Mouseswissprot_no : | <a href="#">Q00196</a>   |
| Immunogen :         | The antiserum was produced against synthesized peptide derived from human OCT2. AA range:1-50  |
| Specificity :       | Oct-2 Polyclonal Antibody detects endogenous levels of 40818 protein.  |
| Formulation :       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source :            | Rabbit   |
| Dilution :          | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.   |
| Purification :      | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| Concentration :     | 1 mg/ml  |
| Storage_stability : | -20°C/1 year   |
| Msds :              | MSDS_Antibody.pdf  |
| Other_name :        | POU2F2; OCT2; OTF2; POU domain; class 2, transcription factor 2; Lymphoid-restricted immunoglobulin octamer-binding protein NF-A2; Octamer-binding protein 2; Oct-2; Octamer-binding transcription factor 2; OTF-2 |




Molecular  
Weight :

52KD

---

## Product Images



Western Blot analysis of various cells using Oct-2 Polyclonal Antibody