

Rag A/B Polyclonal Antibody


| | |
|---------------------|---|
| Catalog_no : | AT3990 |
| Applications : | WB,IHC-p,ELISA |
| Reactivity : | Human,Mouse,Rat |
| Category : | 抗原抗体 |
| Size : | 100µg/50µg/20µg |
| Gene_name : | RRAGA/RRAGB |
| Protein_name : | Ras-related GTP-binding protein A/B |
| Humangene_id : | 10670/10325 |
| Humanswissprot_no : | Q7L523/Q5VZM2 |
| Mousegene_id : | 68441/245670 |
| Ratgene_id : | 117044/100909655 |
| Ratswissprot_no : | Q63486/Q63487 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human RRAGA/B. AA range:264-313 |
| Specificity : | Rag A/B Polyclonal Antibody detects endogenous levels of Rag A/B protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Rabbit |
| Dilution : | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage_stability : | -20°C/1 year |
| Msds : | MSDS_Antibody.pdf |
| Other_name : | RRAGA; Ras-related GTP-binding protein A; Rag A; RagA; Adenovirus E3 14.7 kDa-interacting protein 1; FIP-1; RRAGB; Ras-related GTP-binding protein B; Rag B; RagB |




Molecular
Weight :

34KD


Product Images



Western Blot analysis of various cells using Rag A/B Polyclonal Antibody diluted at 1 : 1000



Western Blot analysis of HepG2 cells using Rag A/B Polyclonal Antibody diluted at 1 : 1000



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

